

# Markscheme

**May 2025**

**Chemistry**

**Standard level**

**Paper 2**

© International Baccalaureate Organization 2025

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2025

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2025

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.


En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.


## Subject Details: Chemistry standard level Paper 2 Markscheme

Candidates are required to answer **ALL** questions. Maximum total = **[50 marks]**.

1. Each row in the “Question” column relates to the smallest subpart of the question.
2. The maximum mark for each question subpart is indicated in the “Total” column.
3. Each marking point in the “Answers” column is shown by means of a tick (✓) at the end of the marking point.
4. A question subpart may have more marking points than the total allows. This will be indicated by “**max**” written after the mark in the “Total” column. The related rubric, if necessary, will be outlined in the “Notes” column.
5. An alternative word is indicated in the “Answers” column by a slash (/). Either word can be accepted.
6. An alternative answer is indicated in the “Answers” column by “**OR**”. Either answer can be accepted.
7. An alternative markscheme is indicated in the “Answers” column under heading **ALTERNATIVE 1** etc. Either alternative can be accepted.
8. Words inside chevrons « » in the “Answers” column are not necessary to gain the mark.
9. Words that are underlined are essential for the mark.
10. The order of marking points does not have to be as in the “Answers” column, unless stated otherwise in the “Notes” column.
11. If the candidate’s answer has the same “meaning” or can be clearly interpreted as being of equivalent significance, detail and validity as that in the “Answers” column then award the mark. Where this point is considered to be particularly relevant in a question it is emphasized by **OWTTE** (or words to that effect) in the “Notes” column.
12. Remember that many candidates are writing in a second language. Effective communication is more important than grammatical accuracy.
13. Occasionally, a part of a question may require an answer that is required for subsequent marking points. If an error is made in the first marking point then it should be penalized. However, if the incorrect answer is used correctly in subsequent marking points then **follow through** marks should be awarded. When marking, indicate this by adding **ECF** (error carried forward) on the script.
14. Do **not** penalize candidates for errors in units or significant figures, **unless** it is specifically referred to in the “Notes” column.
15. If a question specifically asks for the name of a substance, do not award a mark for a correct formula unless directed otherwise in the “Notes” column. Similarly, if the formula is specifically asked for, do not award a mark for a correct name unless directed otherwise in the “Notes” column.
16. If a question asks for an equation for a reaction, a balanced symbol equation is usually expected, do not award a mark for a word equation or an unbalanced equation unless directed otherwise in the “Notes” column.
17. Ignore missing or incorrect state symbols in an equation unless directed otherwise in the “Notes” column.


| Question |     | Answers   | Notes   | Total |
|----------|-----|---|---|-------|
| 1.       | (a) | <p><i>Molecular formula:</i> C<sub>20</sub>H<sub>14</sub>O<sub>4</sub> ✓</p> <p><i>Empirical formula:</i> C<sub>10</sub>H<sub>7</sub>O<sub>2</sub> ✓</p>  |   | 2     |
| 1.       | (b) | <p>C<sub>6</sub>H<sub>5</sub>OH + OH<sup>-</sup> → C<sub>6</sub>H<sub>5</sub>O<sup>-</sup> + H<sub>2</sub>O<br/> <b>OR</b><br/>                     C<sub>6</sub>H<sub>5</sub>OH + H<sub>2</sub>O ⇌ C<sub>6</sub>H<sub>5</sub>O<sup>-</sup> + H<sub>3</sub>O<sup>+</sup><br/> <b>OR</b><br/>                     C<sub>6</sub>H<sub>5</sub>OH ⇌ C<sub>6</sub>H<sub>5</sub>O<sup>-</sup> + H<sup>+</sup> ✓</p> |   | 1     |
| 2.       |     | <p>liquids flow / shape not fixed <b>AND</b> molecules/particles free to move ✓</p> <p>non-compressible / fixed volume <b>AND</b> molecules/particles are close/touching/strongly attracted to each other ✓</p>   | <i>Award [1] for two correct macroscopic properties.</i>  | 2     |
| 3.       | (a) | <p>[N<sub>2</sub>] = 0.545 <b>AND</b> [H<sub>2</sub>] = 0.727 <b>AND</b> [NH<sub>3</sub>] = 0.112 «mol dm<sup>-3</sup>» ✓</p> <p><math>K = \frac{[NH_3]^2}{[N_2][H_2]^3}</math> ✓.</p> <p>«<math>K = \frac{0.112^2}{0.545 \times 0.727^3} \Rightarrow 0.0599</math> ✓.</p>  | <p><i>Award [3] for correct final answer.</i></p> <p><i>Award [3] for <math>K_p=1.66 \times 10^{-9}</math>.</i></p> | 3     |

| Question                       | Answers  | Notes   | Total           |
|--------------------------------|--|---|-----------------|
| <p><b>3.</b>    <b>(b)</b></p> |  <p>sharp increase at time volume of container decreased ✓</p> <p>gradual decrease to second reference point ✓</p> <p>final constant concentration above initial concentration ✓</p> | <p><i>Accept straight line or curve for M2.</i></p>   | <p><b>3</b></p> |
| <p><b>4.</b>    <b>(a)</b></p> | <p>3-bromo-2,2-dimethylbutane ✓</p>  | <p><i>Accept<br/>2-bromo-3,3-dimethylbutane or<br/>2,2-dimethyl-3-bromobutane or 3,3-dimethyl-2-bromobutane.</i></p>  | <p><b>1</b></p> |
| <p><b>4.</b>    <b>(b)</b></p> | <p>«molar mass =&gt;» 165.09 ✓</p> <p>«% H = 13.13 x 100/ 165.09 =&gt;» 7.953% ✓</p>   | <p><i>Award M2 only if answer has four significant figures.</i></p> <p><i>Award [2] for correct final answer.</i></p> | <p><b>2</b></p> |

| Question |     |       | Answers   | Notes  | Total |
|----------|-----|-------|---|--|-------|
| 4.       | (c) |       | any structural isomer of $CH_3CHBrC(CH_3)_3$ . ✓  |  | 1     |
| 4.       | (d) | (i)   |  <p>curly arrow from lone pair/negative charge on O in <math>OH^-</math> to C attached to Br ✓</p> <p>curly arrow from C–Br bond to Br ✓</p>   | <p><i>Ignore bond connectivity and errors in the structure.</i></p> <p><i>Accept S<sub>N</sub>1 mechanism.</i></p> | 2     |
| 4.       | (d) | (ii)  | <p><i>Homolytic fission:</i><br/>each atom receives one «bonding» electron «when bond breaks»<br/><b>OR</b><br/>generates «neutral» free radicals ✓</p> <p><i>Heterolytic fission:</i><br/>one atom receives both «bonding» electrons «when bond breaks»<br/><b>OR</b><br/>generates «charged» ions ✓</p> | <p><i>Award [1 max] if correct descriptions are reversed.</i></p>  | 2     |
| 4.       | (d) | (iii) | <p><i>Any 3 of the following:</i></p> <p>temperature increases kinetic energy/speed of molecules ✓</p> <p>more frequent collisions ✓</p> <p>more molecules have <math>E \geq E_a</math> at higher temperature ✓</p> <p>larger ratio/percentage of collisions are successful ✓</p>                         | <p><i>M2 requires time reference for probability, chance, or number of collisions.</i></p>                         | 3 max |

| Question |     |      | Answers   | Notes  | Total |
|----------|-----|------|---|--|-------|
| 5.       | (a) |      | hydrogen is the limiting reactant <b>OR</b> $2\text{H}_2:1\text{C}_2\text{H}_6$ ✓<br>80.0 cm <sup>3</sup> ✓   | Award <b>[2]</b> for correct final answer.<br>Accept answers expressed in dm <sup>3</sup> .  | 2     |
| 5.       | (b) |      | 60.0 cm <sup>3</sup> of C <sub>2</sub> H <sub>2</sub> in excess/leftover ✓<br>«60.0 cm <sup>3</sup> of C <sub>2</sub> H <sub>2</sub> + 80.0 cm <sup>3</sup> of C <sub>2</sub> H <sub>6</sub> ⇒» 140.0 cm <sup>3</sup> ✓ | Award <b>[2]</b> for correct final answer.<br>Accept answers expressed in dm <sup>3</sup> .  | 2     |
| 6.       | (a) | (i)  | $\text{Li} \rightarrow \text{Li}^+ + \text{e}^-$<br><b>OR</b><br>$\text{LiC}_6 \rightarrow \text{C}_6 + \text{Li}^+ + \text{e}^-$ ✓   | Do <b>not</b> accept 6C.<br>Do <b>not</b> penalize equilibrium arrows.   | 1     |
| 6.       | (a) | (ii) | arrow from cathode to anode in the external circuit ✓   | Do <b>not</b> award mark if any arrows are in the electrolyte.   | 1     |
| 6.       | (b) |      | «battery» stores «surplus» energy «from solar panel» ✓  | Accept “stores charge/power for night use”.<br>Accept “charges battery”.   | 1     |
| 7.       | (a) | (i)  | $1s^2 2s^2 2p^6 3s^2 3p^4$<br><b>OR</b><br>[Ne] $3s^2 3p^4$ ✓   | Accept correct electron configurations with values that are not written with superscripts, e.g. 1s2 2s2 2p6, 1s <sub>2</sub> 2s <sub>2</sub> 2p <sub>6</sub> . | 1     |

| Question |     |       | Answers  | Notes  | Total |
|----------|-----|-------|--|--|-------|
| 7.       | (a) | (ii)  | $\begin{array}{c} \ddot{\text{O}} \\ \vdots \\ \text{:}\ddot{\text{O}}\text{---}\ddot{\text{S}}\text{=}\ddot{\text{O}} \\ \vdots \\ \ddot{\text{O}} \end{array}$ <p>OR</p> $\begin{array}{c} \ddot{\text{O}} \\ \vdots \\ \ddot{\text{O}}\text{=}\ddot{\text{S}}\text{=}\ddot{\text{O}} \\ \vdots \\ \ddot{\text{O}} \end{array} \checkmark$ | <p>Accept the Lewis formulas with the formal charges.</p> <p>Accept any combination of dots or crosses to represent electrons, or lines to represent electron pairs.</p>   | 1     |
| 7.       | (a) | (iii) | <p>Electron domain geometry:<br/>trigonal planar ✓</p> <p>Molecular domain geometry:<br/>bent / V-shaped / angular ✓</p>   | <p>Accept triangular planar for M1.</p> <p>Apply ECF from Lewis formula.</p> <p>Award [1 max] if correct answers are reversed.</p>   | 2     |
| 7.       | (a) | (iv)  | <p>«both» bonds are polar / electronegativity difference «between O and S» ✓</p> <p>«bond» dipoles do not cancel each other / there is a net dipole «because bonds are at an angle less than 180° » ✓</p>  | <p>Accept unsymmetrical distribution of charge <b>OR</b> dipoles add to give a «partial» positive «charge» on S and a «partial» negative «charge» on the O atoms for M2.</p> <p>Apply ECF from molecular geometry.</p>                       | 2     |
| 7.       | (b) | (i)   | <p><math>2\text{SO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{SO}_4(\text{aq})</math><br/>correct product <b>AND</b> state symbols for reactants and product ✓</p> <p>correct balancing ✓</p>  | <p>Accept <math>2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{SO}_3(\text{g})</math><br/><b>AND</b> <math>2\text{SO}_3(\text{g}) + 2\text{H}_2\text{O}(\text{l}) \rightarrow 2\text{H}_2\text{SO}_4(\text{aq})</math></p> | 2     |
| 7.       | (b) | (ii)  | <p>sulfur oxidation state changed from «+»4 to «+»6 ✓</p> <p>oxygen «atoms in O<sub>2</sub>» oxidation state changed from 0 to -2 ✓</p>  | <p>Accept oxidation state of S increased by 2 OR oxygen reduced by 2 for [1].</p> <p>Accept increase from IV to VI.</p>  | 2     |

| Question |     |       | Answers   | Notes  | Total |
|----------|-----|-------|---|--|-------|
| 7.       | (b) | (iii) | limestone/building dissolves ✓<br>$\text{CaCO}_3(\text{s}) + \text{H}_2\text{SO}_4(\text{aq}) \rightarrow \text{CaSO}_4(\text{aq}) + \text{H}_2\text{O}(\text{l}) + \text{CO}_2(\text{g}) \checkmark$                             | Accept limestone becomes rough/loss of carved details/edges are rounded/erodes/OWTTE for M1.<br>Accept $\text{H}_2\text{CO}_3$ product.  | 2     |
| 7.       | (c) |       | Any one of:<br>  | At least one correct bond required.<br>Accept any type of dotted line.<br>Do <b>not</b> accept an intramolecular hydrogen bond.<br>Do <b>not</b> accept any other bonds (eg H--H)<br>Do <b>not</b> accept multiple bonds from a single hydrogen. | 1     |
| 7.       | (d) |       | covalent bonds between silicon atoms ✓<br>London/dispersion forces between poly(ethene) «chains/molecules» ✓<br>covalent bonds «much» stronger than London/dispersion/intermolecular forces «hence Si has higher melting point» ✓ | M3: Do <b>not</b> accept “bonding in silicon is stronger than poly(ethene)” without named bonding/forces.  | 3     |
| 8.       | (a) | (i)   | $6\text{CO}_2(\text{g}) + 6\text{H}_2\text{O}(\text{l}) \rightarrow \text{C}_6\text{H}_{12}\text{O}_6(\text{aq}) + 6\text{O}_2(\text{g}) \checkmark$  |  | 1     |

| Question |     |      | Answers  | Notes  | Total |
|----------|-----|------|--|--|-------|
| 8.       | (a) | (ii) | <p>Any two:</p> <p>«ethanol is» renewable / sustainable resource ✓</p> <p>«ethanol has» low/zero carbon footprint / produces less CO<sub>2</sub> ✓</p> <p>less sulfur dioxide «than fossil fuels»<br/> <b>OR</b><br/>                     less acid rain «than fossil fuels» ✓</p> <p>less incomplete combustion «than fossil fuels»<br/> <b>OR</b><br/>                     less carbon monoxide/soot «than fossil fuels» ✓</p> | <p>Accept “ethanol is biodegradable /less toxic than gasoline”.</p> <p>Do <b>not</b> accept just “less harmful”.</p> | 2 max |
| 8.       | (b) |      | <p>«<math>Q = (4.00 / 46.08) \times 1367 \Rightarrow 119 \text{ kJ} / 119000 \text{ J}</math>» ✓</p> <p>«<math>\Delta T = Q / mc = 119000 / (500.0 \times 4.18) \Rightarrow 56.9 \text{ «K»}</math>» ✓</p>   | Accept 56.8 «K».   | 2     |